arXiv:gr-qc/0610030v2 15 Apr 2007

Geometric Flows and Black Holes

Fu-Wen Shu[1, 2, 3, *] and You-Gen Shen[1,3, 4, †]

[1]*Shanghai Astronomical Observatory, Chinese Academy of Sciences, Shanghai 200030, P. R. China*
[2]*Graduate School of Chinese Academy of Sciences, Beijing 100039, P. R. China*
[3]*Joint Institute for Galaxy and Cosmology of SHAO and USTC, Shanghai 200030, P. R. China*
[4]*National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100012, P. R. China*

(Dated: February 3, 2008)

Motivated by the newest progress in geometric flows both in mathematics and physics, we apply the geometric evolution equation to study some black-hole problems. Our results show that, under certain conditions, the geometric evolution equations satisfy the Birkhoff theorem, and surprisingly, in the case of spherically symmetric metric field, the Einstein equation, the Ricci flow, and the hyperbolic geometric flow in vacuum spacetime have the same black-hole solutions, especially in the case of $\Lambda = 0$, they all have the Schwarzschild solution. In addition, these results can be generalized to a kind of more general geometric flow.

PACS numbers: 02.30.Jr, 02.40.Ky, 04.20.Jb, 04.70.-s, 04.70.Bw

Introduction. The geometric flows provide new ways to address a variety of non-linear problems in Riemannian geometry. In addition, it is shown that they also play an important role in physics, especially Einstein equation(it is included as the limiting case of no flow) in general relativity, and Ricci flow [1] in the renormalization group analysis of non-linear sigma models in string theory[2, 3] and various models of boundary quantum field theory of current interest[4]. Recently, Graf [5] showed that the Einstein gravity is a limiting case of the Ricci flow gravity. After that, Headrick and Wiseman [6] found that the Ricci flow can be used to study black holes. As to other things, Perelman [7, 8] applied the Ricci flow to prove Thurston's geometrization conjecture [9] concerning the classification of 3-manifolds (for reviews see [10]).

Recently Kong and Liu [11] introduced a kind of new flow, which they called the *hyperbolic geometric flow*. It is a system of non-linear evolution partial differential equations of second order. In their remarkable work on hyperbolic flow [11], they showed that "*the hyperbolic geometric flow is a natural and powerful tool to study some problems arising form differential geometry such as singularities, existence and regularity*".

Motivated by these amazing progress of the geometric flows both in mathematics and physics, we study the gravitational properties of the geometric evolution equations. Thanks to recent introduction of the hyperbolic geometric flow, we have a set of general evolution equations including the Einstein equations (no flow case), the Ricci flow, and the hyperbolic flow. These equations are [11]

$$\alpha_{\mu\nu}\frac{\partial^2 g_{\mu\nu}}{\partial t^2} + \beta_{\mu\nu}\frac{\partial g_{\mu\nu}}{\partial t} + \gamma_{\mu\nu} g_{\mu\nu} + 2R_{\mu\nu} = 0. \qquad (1)$$

where $\alpha_{\mu\nu}$, $\beta_{\mu\nu}$, $\gamma_{\mu\nu}$ are certain smooth functions (may depend on t) on n–dimensional complete Riemannian manifold $\mathcal{M}$, and $g_{\mu\nu}$ is Riemannian metric, $R_{\mu\nu}$ is Ricci tensor. Obviously, in the case of $\alpha_{\mu\nu} = \beta_{\mu\nu} = 0$ and $\gamma_{\mu\nu} = constant$, Eq. (1) corresponds to the Einstein equation with regard to vacuum spacetime; the case of $\alpha_{\mu\nu} = 0$, $\beta_{\mu\nu} = 1$ and $\gamma_{\mu\nu} = 0$ corresponds to the Ricci flow, which is analogous to the heat equation in physics; and the case of $\alpha_{\mu\nu} = 1$, $\beta_{\mu\nu} = \gamma_{\mu\nu} = 0$ corresponds to the hyperbolic geometric flow, which is analogous to the wave equation in physics, describing the wave character of the metrics and curvatures of manifolds.

This letter is an attempt to apply the geometric evolution equation (1) to study some black-hole problems. Our results show that, under certain conditions, the Birkhoff theorem[15] [12] is still reserved in the frame of the geometric flow discussed here. We also show that for the spherically symmetric metric in the 4–dimensional Riemannian manifold, the Einstein equation, the Ricci flow, and the hyperbolic geometric flow in vacuum spacetime all have the same black-hole solutions. In addition, we can generalize our results to a kind of more general geometric flow.

Birkhoff theorem. In 1923, Birkhoff [12, 14] showed that any spherically symmetric solution of the vacuum Einstein field equations must be static and asymptotically flat. In this section we will try to extend the Birkhoff theorem to the case including geometric evolution (1).

Theorem 1. *For the geometric flow (1) in the 4–dimensional Riemannian manifold, if $\alpha_{\mu\nu}$, $\beta_{\mu\nu}$, and $\gamma_{\mu\nu}$ are independent of t and r, then Eq. (1) satisfies the Birkhoff theorem.*

Proof. Consider a line element with spherically symmetric metric

$$g_{\mu\nu} = diag\left(B(r,t), -A(r,t), -r^2, -r^2\sin^2\theta\right),$$

where " *diag* " represents a diagonal matrix. Then we get,

$$\begin{cases} R_{rr} = \frac{B''}{2B} - \frac{B'^2}{4B^2} - \frac{A'B'}{4AB} - \frac{A'}{Ar} - \frac{\ddot{A}}{2B} + \frac{\dot{A}\dot{B}}{4B^2} + \frac{\dot{A}^2}{4AB}, \\ R_{\theta\theta} = -1 + \frac{1}{A} - \frac{rA'}{2A^2} + \frac{rB'}{2AB}, \\ R_{\varphi\varphi} = R_{\theta\theta} \sin^2\theta, \\ R_{tt} = -\frac{B''}{2A} + \frac{A'B'}{4A^2} - \frac{B'}{Ar} + \frac{B'^2}{4AB} + \frac{\ddot{A}}{2A} - \frac{\dot{A}\dot{B}}{4AB} - \frac{\dot{A}^2}{4A^2}, \\ R_{rt} = -\frac{\dot{A}}{Ar}, \end{cases} \tag{2}$$

where “ $'$ ” and “ $\cdot$ ” represent derivatives with respect to r and t, respectively. On account of the fact that $g_{\mu\nu}$ is diagonal, from (1) we directly have

$$R_{rt} = \dot{A} = 0, \tag{3}$$

From here we learn that $A(r,t)$ is independent of t, so we let $A(r,t) \equiv a(r)$. Plugging (3) into formula (2), and letting $B(r,t) \equiv \rho(t)b(r)$, from (1) one has

$$\begin{cases} \alpha_{00}\ddot{\rho}b + \beta_{00}\dot{\rho}b + \gamma_{00}\rho b = -2\rho\left(-\frac{b''}{2a} + \frac{a'b'}{4a^2} - \frac{b'}{ar} + \frac{b'^2}{4ab}\right), \\ \gamma_{11}a = -2\left(\frac{b''}{2b} - \frac{b'^2}{4b^2} - \frac{a'b'}{4ab} - \frac{a'}{ar}\right), \\ \gamma_{22}r^2 = -2\left(-1 + \frac{1}{a} - \frac{ra'}{2a^2} + \frac{rb'}{2ab}\right). \end{cases} \tag{4}$$

The first formula in (4) can be decoupled as

$$\begin{aligned} \lambda &= \alpha_{00}\frac{\ddot{\rho}}{\rho} + \beta_{00}\frac{\dot{\rho}}{\rho} + \gamma_{00} && (5) \\ &= -2\left(-\frac{b''}{2ab} + \frac{a'b'}{4a^2b} - \frac{b'}{abr} + \frac{b'^2}{4ab^2}\right), && (6) \end{aligned}$$

where λ is a constant. From (5), one can easily obtain the explicit expression of $\rho(t)$. In the end, we can rescale the time coordinate as

$$dt'^2 \equiv \rho(t)dt^2.$$

In this way, we obtain

$$ds^2 = b(r)dt'^2 + a(r)dr^2 + r^2d\Omega^2,$$

with $a(r)$ and $b(r)$ independent of t. This leads to the Birkhoff theorem in general relativity.

In fact, from the mathematical point of view, we can extend Eq. (1) to the following more general evolution equations

$$\begin{aligned} &\alpha^n_{\mu\nu}\frac{\partial^n g_{\mu\nu}}{\partial t^n} + \alpha^{n-1}_{\mu\nu}\frac{\partial^{n-1} g_{\mu\nu}}{\partial t^{n-1}} + \cdots \\ &+ \alpha^1_{\mu\nu}\frac{\partial g_{\mu\nu}}{\partial t} + \alpha^0_{\mu\nu}g_{\mu\nu} + 2R_{\mu\nu} = 0, \end{aligned} \tag{7}$$

where $\alpha^n_{\mu\nu}$ are certain smooth functions (may depend on t) on m–dimensional complete Riemannian manifold $\mathcal{M}$, and n is integer. Note that for the case of $n > 2$, we have little knowledge of its physical counterpart of Eq. (7). Similarly, one can easily obtain a more general theorem:

Theorem 2. *For the geometric flow (7) in the 4–dimensional Riemannian manifold, if $\alpha^n_{\mu\nu}$ are independent of t and r, then Eq. (7) satisfies the Birkhoff theorem.*

One can prove Theorem 2 by using the same technique addressed above for Theorem 1.

Exact black-hole solutions of geometric evolution equation. As we have learned in general relativity, if the matter distributes spherically inside a ball, and the outside of the ball is vacuum, then we have an exact solution of the Einstein field equation, which is now called Schwarzschild solution. In this section, we will show that not only the Einstein equation has black-hole solution, the Ricci evolution equation and the hyperbolic evolution equation also have. In order to see this in detail, we let $\gamma_{\mu\nu} = 0$ in Eq. (1), so that it corresponds to Ricci flow as $\alpha_{\mu\nu} = 0$, $\beta_{\mu\nu} = 1$, and hyperbolic flow as $\alpha_{\mu\nu} = 1$, $\beta_{\mu\nu} = 0$, respectively. In this case, when taking Eq. (6) into account, Eq. (4) becomes

$$\begin{cases} -\frac{b''}{2ab} + \frac{a'b'}{4a^2b} - \frac{b'}{abr} + \frac{b'^2}{4ab^2} = -\frac{1}{2}\lambda, \\ \frac{b''}{2b} - \frac{b'^2}{4b^2} - \frac{a'b'}{4ab} - \frac{a'}{ar} = 0, \\ -1 + \frac{1}{a} - \frac{ra'}{2a^2} + \frac{rb'}{2ab} = 0. \end{cases} \tag{8}$$

Solving it, one gets

$$y'r + y - 1 + \Lambda r^2 = 0, \tag{9}$$

where $y \equiv \frac{1}{a}$ and $\Lambda = \frac{\lambda}{4}$.

Integrating Eq. (9), we have

$$y = 1 - \frac{2m}{r} - \frac{1}{3}\Lambda r^2,$$

where m is an integration constant (one can easily confirm that $m = \frac{GM}{c^2} > 0$ is the mass of the Schwarzschild black hole). Consequently, we obtain

$$\begin{aligned} a &= \left(1 - \frac{2m}{r} - \frac{1}{3}\Lambda r^2\right)^{-1}, && (10) \\ \frac{b'}{b} &= 2\Lambda ar - \frac{a'}{a}. && (11) \end{aligned}$$

Case 1 $\Lambda = 0$: In this case, integrating Eq.(11) directly, one has $b = \frac{\varepsilon}{a}$, where ε is an integration constant which can be set to 1. In the end, one has

$$ds^2 = \left(1 - \frac{2m}{r}\right)c^2dt^2 - \left(1 - \frac{2m}{r}\right)^{-1}dr^2 + r^2d\Omega^2.$$

This is just the Schwarzschild metric!

Case 2 $\Lambda \neq 0$: In this case, we rewrite $\frac{1}{a}$ as

$$1 - \frac{2m}{r} - \frac{\Lambda}{3}r^2 = -\frac{\Lambda(r - r_a)(r - r_b)(r - r_c)}{3r},$$

where r_a, r_b and r_c are three roots of $\frac{1}{a} = 0$, which depend on the value of the decoupled constant Λ. Plugging this formula into Eq.(11), and integrating it, we have

$$b = \varepsilon(r - r_a)^{\sigma_a}(r - r_b)^{\sigma_b}(r - r_c)^{\sigma_c} \cdot \frac{1}{a},$$

where ε is an integration constant which can be set to 1, and

$$\sigma_i = \frac{\Lambda r_i}{\kappa_i},$$

where $\kappa_i \equiv \frac{1}{2}\frac{d(1/a)}{dr}\mid_{r=r_i}$, which corresponds to the surface gravity of the black hole. In the end, we obtain

$$\begin{aligned} ds^2 &= u(r)\left(1-\frac{2m}{r}-\frac{\Lambda}{3}r^2\right)c^2dt^2 \\ &- \left(1-\frac{2m}{r}-\frac{\Lambda}{3}r^2\right)^{-1}dr^2 + r^2 d\Omega^2, \end{aligned}$$

where

$$u(r) = (r-r_a)^{\sigma_a}(r-r_b)^{\sigma_b}(r-r_c)^{\sigma_c}.$$

(a) $\frac{1}{\Lambda} \geq \frac{9}{4}r_g^2$, where $r_g = 2m$ is the Schwarzschild radius: In this case, r_a, r_b, r_c are real. Let $r_a < r_b < r_c$. After some mathematical analyses, one finds that $r_a < 0$ and $r_c \geq r_b > 0$. Consequently, black hole in this case has two horizons, one of them is the event horizon.

(b) $\frac{1}{\Lambda} < \frac{9}{4}r_g^2$: The black hole only has one horizon—the event horizon (here we let r_a be the event horizon). $r_b = r_c^*$ are complex in this case.

Surprisingly, as we solve the exact solutions of the more general geometric flows (7), we find that we get the completely same solutions as done above. This implies that *any 4–dimensional spherically symmetric metric in vacuum spacetime will inevitably form static black holes, as long as it keeps unchanged or changes isotropically.* This leads to a conjecture, namely, *any geometric flow which satisfies the Birkhoff theorem may have a black-hole solution.*

Conclusions. We have shown that for the geometric flows (1) in the 4–dimensional Riemannian manifold, including the Einstein equations(no flow), the Ricci flow, and the hyperbolic geometric flow, if $\alpha_{\mu\nu}$, $\beta_{\mu\nu}$, and $\gamma_{\mu\nu}$ are independent on t and r, then Eq. (1) satisfies the Birkhoff theorem, namely, any spherically symmetric solution of the vacuum geometric evolution equations must be static and asymptotically flat. Moreover, we can extend the Birkhoff theorem to the more general flow (7), under the condition that $\alpha^n_{\mu\nu}$ are independent of r and t. This implies that *the spherically symmetric change of the initial spherically symmetric metric in the vacuum spacetime does not induce pure gravitational radiation, regardless of the way of this change.*

As to the exact solution of the geometric evolution equations, our results have shown that in the 4–dimensional case of matter distributed spherically inside a ball, and vacuum outside of the ball, the Einstein equation, the Ricci flow, and the hyperbolic geometric flow in vacuum spacetime have the same black-hole solution. Particularly, in the case of $\Lambda = 0$, they all give the Schwarzschild solution.

Acknowledgments. F. W. Shu wishes to thank Kefeng Liu for informing me about the reference [11]. The work was supported by the National Natural Science Foundation of China under Grant No. 10573027 and 10663001, and the Foundation of Shanghai Natural Science Foundation under Grant No. 05ZR14138.

---

[*] Electronic address: fwsu@shao.ac.cn

[†] Electronic address: ygshen@shao.ac.cn

[1] R. Hamilton, J. Differential Geom. **17** 255-306 (1982).

[2] D. H. Friedan, Ann. Phys. **163** 318 (1985).

[3] R. Leigh, Mod. Phys. Lett. A **4** 2767 (1989).

[4] S. Lukyanov, E. Vitchev and A. Zamolodchikov, Nucl. Phys. B **683** 423 (2004).

[5] W. Graf, arXiv: gr-qc/0602054.

[6] M. Headrick and T. Wiseman arXiv: hep-th/0606086.

[7] G. Perelman, arXiv: math. DG/0211159.

[8] G. Perelman, arXiv: math. DG/0303109.

[9] W. P. Thurston, Bull. Amer. Math. Soc. **6** 357 (1982).

[10] J. W. Morgan, Bull. Amer. Math. Soc. (N.S.) **42** 57 (2005).

[11] D. X. Kong and K. Liu, http://www.cms.zju.edu.cn/UploadFiles/AttachFiles/2006828859465

[12] G. D. Birkhoff, *Relativity and modern physics*, Cambridge: Cambridge university press, (1923).

[13] S. Deser, Gen. Relativ. Gravit. **37** 2251 (2005).

[14] J. T. Jebsen, Gen. Relativ. Gravit. **37** 2253 (2005).

[15] In fact, recently Deser[13] pointed out that this theorem was first discovered by Jebsen [14], instead of Birkhoff.